

5D Kaluza - Klein's Wormhole Between Two Event Horizons

Dzhunushaliev V.D. *

Theoretical Physics Department

Kyrgyz State National University, Bishkek, 720024

The solution of the 5D Kaluza-Klein's theory is obtained. This solution is a Lorentzian wormhole between two event horizons. It is shown that this solution can sew with (4D Reissner-Nordström's solution + Maxwell's electrical field) on the event horizon. Such construction is composite wormhole connecting two asymptotically flat region. From the viewpoint of infinite observer this wormhole is an electrical charge. According to J.Wheeler terminology this is "charge without charge".

04.50.+h

In contemporary models of quantum gravity it is of prime important the dynamical process carrying out change of the space - time topology [1]. It is supposed that such processes take place as a result of the wormholes origin. There are many models of the Euclidean's wormholes filled some kind of matter field (see, for example, [2], [3]).

The aim of this paper is the construction of a composite wormhole: it is the solution of 5D Einstein's equations (Kaluza - Klein's equation) under event horizon (EH), and Reissner - Nordström's solution outside EH. Onto EH is performed sewing the 5D Kaluza - Klein's metric with 4D Einstein's metric + electrical field.

A. 5D Wormhole in Kaluza - Klein's theory

The initial 5D Kaluza - Klein's equation's write down in the following standart manner:

$$R_{\mu\nu} - \frac{1}{2}G_{\mu\nu}R = 0 \quad (1)$$

where $G_{\mu,\nu}$ is 5D metrical tensor; $R_{\mu\nu}$ is Ricci tensor; R is scalar curvature; $\mu, \nu = 1, 2, 3, 4, 5$. 5D metric we seek in the following wormholelike view:

$$ds^2 = e^{2\nu(r)}dt^2 - e^{2\psi(r)}(d\chi - \omega(r)dt)^2 - dr^2 - a^2(r)(d\theta^2 + \sin^2\theta d\varphi^2), \quad (2)$$

where χ is 5 supplementary coordinate; r, θ, φ are 3D polar coordinates; t is the time. Corresponding Kaluza - Klein's equations have the following view:

$$\psi'' + \psi'^2 + \psi'\nu' + \frac{2a'\psi'}{a} + \frac{q^2}{2a^4}e^{-4\psi} = 0, \quad (3)$$

$$a'' + a'\psi' + \frac{a'^2}{a} + a'\nu' - \frac{1}{a} = 0, \quad (4)$$

$$\nu'' + \nu'^2 + \psi'\nu' + \frac{2a'\nu'}{a} - \frac{q^2}{2a^4}e^{-4\psi} = 0, \quad (5)$$

$$\psi'' + \frac{2a''}{a} + \psi'^2 - \psi'\nu' - \frac{2a'\nu'}{a} = 0, \quad (6)$$

here $(')$ denotes the derivative with respect to r . In equations system (3–6) it is taken the following equation into account:

$$\omega' = \frac{q}{a^2}e^{-3\psi+\nu}, \quad (7)$$

obtained by integration of $R_{\chi t} = 0$ equation. q in Eq.(7) is some constant ("electrical" Kaluza - Klein's charge).

From Eq's (3) and 5 it follows that $\psi = -\nu$ and the initial system (3–6) can written as follows:

$$a'' + \frac{a'^2}{a} - \frac{1}{a} = 0, \quad (8)$$

$$\nu'' + \frac{2a'\nu'}{a} - \frac{q^2}{2a^4}e^{4\nu} = 0, \quad (9)$$

$$\nu'^2 + \frac{1}{a^2} \left(1 - \frac{q^2}{4a^2}e^{4\nu} \right) - \frac{a'^2}{a^2} = 0. \quad (10)$$

The integration of the Eq.(8) and Eq.(10) leads to the following result:

$$a^2 = r_0^2 + r^2, \quad (11)$$

$$e^{2\nu} = \frac{2r_0}{q} \frac{r_0^2 + r^2}{r^2 - r_0^2}. \quad (12)$$

where r_0 is a minimal radius of given wormhole. Eq.(9) is identical satisfied.

The integration of Eq.(7) leads to the following result:

$$\omega = \frac{4r_0^2}{q} \frac{r}{r_0^2 - r^2}. \quad (13)$$

It can be shown that the time component of metrical tensor $G_{tt}(r = \pm r_0) = 0$. This indicates that there is the event horizon by $r = \pm r_0$.

It is easy to see that solution (11–13) is nonsingular by $r \in (-\infty, +\infty)$ but the supplementary coordinate (χ) can turns out to the timelike coordinate by $|r| > r_0$. This is unsatisfactory with respect to physical point of view. On the other hand if the Kaluza - Klein's action:

*E-mail address: dzhun@freenet.bishkek.su

$$S = - \int R \sqrt{-G} d^5 x \quad (14)$$

not vary with respect to G_{tt} component, then we obtain the regular physically reasonable solution by $|r| > r_0$. This is Reissner-Nordström's solution.

B. The sewing Kaluza - Klein's wormhole with Reissner - Nordström's solution

In this section we show that the Reissner - Nordström's solution really can to sew on EH with above received 5D wormhole [4]. To do this, the 5D Kaluza - Klein's metrical tensor is necessary to sew with (4D metric + Maxwell's electrical field in Reissner - Nordström's solution).

The following metric components must be in agreement on the sewing surface = EH :

$$e^{2\nu_0} - \omega_0^2 e^{-2\nu_0} = G_{tt}(r_0) = g_{tt}(r_+) = 0, \quad (15)$$

$$r_0^2 = G_{\theta\theta} = g_{\theta\theta} = r_+^2, \quad (16)$$

where G and g are 5D and 4D metrical tensor respectively. $r_+ = m + \sqrt{m^2 + Q^2}$ is EH for Reissner - Nordström's solution (m and Q are mass and charge of the Reissner - Nordström's black hole). The quantity marked by (0) sign are taken by $r = r_0$.

To sew $G_{\chi t}$ and 4D electrical field we consider 5D $R_{\chi t} = 0$ equation and Maxwell's equation:

$$[a^2 (\omega' e^{-4\nu})]' = 0, \quad (17)$$

$$(r^2 E_r)' = 0, \quad (18)$$

here E_r is 4D electrical field.

This 2 equations are practically the Gauss law and they indicate that some quantity multiplied by area is conserved. In 4D case this quantity is 4D Maxwell's electrical field and from this follows that the electrical charge is conserved. Thus, naturally we must join 4D electrical Reissner - Nordström's field $E_{RN} = Q/r_+^2$ with "electrical" Kaluza - Klein's field $E_{KK} = \omega' e^{-4\nu}$ on EH:

$$\omega_0' e^{-4\nu_0} = \frac{q}{2r_0^2} = E_{KK} = E_{RN} = \frac{Q}{r_+^2}. \quad (19)$$

Now, it is necessary to consider the question on the sewing physical quantity produced from 1th and 2th derivative of metric. Such geometrical quantity as connection and curvature tensor consist of the 1th and 2th derivative of the metrical tensor component respectively. Hence, it is necessary to examine the question of sewing this quantity.

By sewing 4D and 5D connections and curvature tensors on EH we obtain a singularity (on surface sewing) similar to singularity of $|x|$ function at the point $x = 0$. In this point of view 1th derivative has discontinuity and 2th derivative has δ -like singularity. Analogically in our

situation the connection has discontinuity and curvature tensor has δ -like singularity.

By such singularity the connection as well as the curvature tensor can to have certain nonsingular geometrical meaning like to gravitational Regge calculus: The components of the tangent vector, which are transferred in a parallel way through sewing surface (EH) at an infinitesimal distance, acquire the finite increment from the fact that the connection has discontinuity.

-
- [1] J.Wheeler, In Relativity Groups and Topology, eds. B.S. and C.M.DeWitt. Gordon and Breach, N.-Y. (1963).
 - [2] S.J.Rey, Nucl.Phys., **v.336**, 146(1990).
 - [3] K.Joshida, S.Hirenzaki, K.Shiraishi, Phys.Rev, **D42**, 1973(1990).
 - [4] V.Dzhunushaliev, Izvestia vuzov, ser.Fizika, N6, 78(1993).